

AD-A105 061 NAVAL UNDERWATER SYSTEMS CENTER NEW LONDON CT NEW LO--ETC F/G 20/1
REGIONAL DEPENDENCE OF VERY LOW-FREQUENCY SOUND ATTENUATION IN --ETC(U)
SEP 81 D G BROWNING, M J FECHER, R H MELLEN
UNCLASSIFIED NUSC-TO-6561 NL

[11F]
ALL
REPROD



				END
				DATE
				FILED
				10 81
				DTIC

11) NUSC Technical Document 6561
22 September 1981

LEVEL II

12

Regional Dependence of
Very Low-Frequency Sound Attenuation
in the Deep Sound Channel: Correlation
with Internal Wave Measurements,

A Paper Presented at the Joint Meeting of the
Acoustical Society of America and the Acoustical
Society of Japan, 27 November--1 December 1978,
Honolulu, Hawaii,

David G. Browning
Michael J. Fecher
Robert H. Mellen
Surface Ship Sonar Department

DTIC
ELECTE
OCT 6 1981
H



Naval Underwater Systems Center
Newport, Rhode Island / New London, Connecticut


Approved for public release; distribution unlimited.

405918 81 10 2 086

Preface

This document was prepared under the sponsorship of the Naval Material Command under NUSC Project No. A65410, "Acoustic Variability Within the Sound Channel," as part of the NUSC Independent Research Program; NAVMAT Program Manager, CAPT D. F. Parrish, and NUSC Principal Investigator, D. G. Browning.

Reviewed and Approved: 22 September 1981


W. A. Von Winkle
Associate Technical Director
for Technology

**The authors of this document are located at the
New London Laboratory, Naval Underwater Systems Center,
New London, Connecticut 06320.**

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER TD 6561	2. GOVT ACCESSION NO. 4D-1105061	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) REGIONAL DEPENDENCE OF VERY LOW-FREQUENCY SOUND ATTENUATION IN THE DEEP SOUND CHANNEL: CORRELATION WITH INTERNAL WAVE MEASUREMENTS, A Paper Presented at the Joint Meeting of the ASA and the ASJ, 27 Nov.-1 Dec. 1978, Honolulu, Hawaii		5. TYPE OF REPORT & PERIOD COVERED
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) David G. Browning, Michael J. Fecher, and Robert H. Mellen		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Underwater Systems Center New London Laboratory New London, CT 06320		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS A65410
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Material Command Washington, DC 20360		12. REPORT DATE 22 September 1981
		13. NUMBER OF PAGES 12
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION / DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Attenuation Internal Waves Scattering		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This document presents the oral and visual presentation entitled "Regional Dependence of Very Low-Frequency Sound Attenuation in the Deep Sound Channel: Correlation with Internal Wave Measurements," presented at the joint meeting of the Acoustical Society of America and the Acoustical Society of Japan, 27 November-1 December 1978, Honolulu, Hawaii. The most promising mechanism for the very low-frequency sound attenuation observed in the deep sound channel is diffusive scattering by internal		

20. (Continued)


waves. Mellen et al. (J. Acoust. Soc. Am., 60, 1053-1056 (1976)) have obtained estimates for the extra attenuation using the Garrett-Munk internal wave model and found consistency with the lower experimental values reported. Kibblewhite et al. (J. Acoust. Soc. Am. 63, 1169-1177 (1978)) have shown a definite regional dependence in the Pacific.

In this paper we compare regional oceanographic measurements with the Garrett-Munk internal wave model and also correlate local acoustic measurements with estimates of the extra attenuation.

Acc.	✓
NE	
DE	
U	
J	
By	
DE	
AN	
Dist	
A	

**Regional Dependence of Very Low-Frequency
Sound Attenuation in the Deep Sound Channel:
Correlation with Internal Wave Measurements**

SOUND ATTENUATION IN THE OCEAN



Slide 1


At most frequencies, measurements of the attenuation of sound in the sea can be explained by chemical absorption mechanisms. For very low frequencies, typically below 100 Hz, the measured attenuation is consistently higher than expected from absorption alone.

This excess attenuation, generally attributed to diffusive scattering loss from oceanographic inhomogeneities, ranges from 4 to 50×10^{-4} dB/km depending upon the geographic location of the experiment.

Mellen, Browning, and Goodman* have estimated the attenuation coefficient for an idealized deep sound channel perturbed by internal waves based on the Garrett-Munk formulation. The result, 5×10^{-4} dB/km, is consistent with the lower experimental values.

The following discussion will show that improved predictions of low frequency attenuation coefficients can be obtained using internal wave theory if regional deviations from the ideal ocean are considered.

*"Diffusion Loss in a Stratified Sound Channel, *Journal of the Acoustical Society of America*, vol. 60, no. 5, November 1976.



Slide 2

For long-range diffusion loss in a stratified sound channel, we consider a plane acoustic wave propagating through random lenticular sound speed inhomogeneities that have large-scale dimensions compared with the acoustic wavelength. In particular, an exponential buoyancy frequency profile N (dashed line) and the resulting canonical sound speed profile C are assumed. Typical values are shown. B is the vertical scale of the sound channel. Internal wave displacements produce inhomogeneities by perturbing the sound speed profile, as suggested on the right-hand side.

TRANSVERSE DIFFUSION CONSTANT AT SOUND CHANNEL AXIS

$$D_1 \approx 4\mu_1^2 j * n_1 \left(\frac{n_1}{\omega_1} \right)$$

ATTENUATION COEFFICIENT FOR DIFFUSIVE SCATTERING

$$\alpha_1 = \frac{13 D_1}{\theta_0^2 B} \text{ dB/km}$$

where,

- μ_1^2 = variance of refractive index $\equiv \overline{\left(\frac{\Delta c}{c} \right)^2}$
- θ_0 = initial angle of bottom-limited array
- B = scale size of sound channel
- n_1 = (scaled) buoyancy frequency
- ω_1 = (scaled) inertial frequency
- $j *$ = internal wave mode scale number, typically = 3

Slide 3

Then, for small ray angles, the vertical transverse diffusion constant and the attenuation coefficient are evaluated at the sound channel axis, as indicated by subscript 1, μ^2 is the variance of refractive index, and θ_0 is the initial angle for the bottom-grazing ray.

Values of μ^2 , B , n , and θ_0 have been inferred from two sets of oceanographic data collected at mid-latitudes in the North Atlantic. Resulting attenuation coefficients will be compared with the acoustically-derived coefficients.

$$\text{GENERAL} \quad \mu^2(z) \equiv \left[\frac{\Delta c(z)}{C} \right]^2 = \left[\frac{\partial_z C_p(z)}{C} \right]^2 \zeta^2(z) \quad (1)$$

INTERNAL WAVE - CANONICAL MODEL

$$(2) \quad \zeta^2(z) = \zeta_o^2 \frac{N_o}{N(z)}, \quad \zeta(z) = \text{rms IW displacement } z=0 = 7.3 \text{ m}$$

$$(3) \quad \frac{\partial_z C_p(z)}{C} = \eta(z) N^2(z) / g \\ \approx \eta_o N_o^2 / g [\exp(-2z/B)]$$

$$\text{where } \eta_o = 24.5 \approx \eta(z)$$

$$(4) \quad N(z) = N_o \exp(-z/B), \quad N_o = 3 \text{ cph}, B = 1 \text{ km}$$

$$(5) \quad \mu_c^2(z) = \mu_o^2 \exp(-3z/B) \approx 2.4 \times 10^{-7} \exp(-3z/B)$$


INTERNAL WAVE - NON-CANONICAL

$$(6) \quad \mu_{nc}^2(z) = \frac{\langle [\partial_z C_p(z)]^2 \rangle}{C^2} \langle \zeta_d^2 \rangle \frac{N_d}{\langle N(z) \rangle}$$

Slide 4


But first we recall that mean-square fluctuations in sound speed (μ^2) are related to vertical perturbations by equation 1. Here, zeta is the rms displacement amplitude, and $\partial_z C_p$ is the potential sound speed gradient. Variations in internal wave amplitude with depth can be scaled according to the buoyancy profile (as in equation 2). The potential sound speed gradient is also related to N by equation 3. Assuming idealized exponential stratification (equation 4) then yields equation 5, a canonical model wherein μ^2 decays exponentially with depth.

Equation 6 is non-canonical in that exponential stratification is not assumed. Instead, the internal waves perturb the actual sound speed profile; wave amplitude is scaled to the existing buoyancy profile and a measured rms displacement at depth d . One set of measurements used to evaluate equation 6 was conducted in the Azores Fixed Acoustic Range.

**Slide 5**

This is a composite of 46 sound speed profiles collected in the Azores Range. Three similar data sets were collected concurrently with profiling systems onboard other ships. Each data set shows the same spread in sound velocities, and the intrusion of the Mediterranean water at 900 meters.

These fluctuations in sound speed were used to calculate μ^2 directly.




Slide 6

Here μ^2 has been scaled as dB relative to 10^{-6} . The thick solid line is the canonical internal wave model. Above 600 meters and below 1000 meters, the measured values agree with the canonical decay rate.

However, at the deep sound axis (~ 1500 meters), the observed μ^2 is at least 17 dB larger than given by the canonical equation.


To infer whether these observed values can be attributed to internal waves in the real ocean, μ^2 was calculated using the non-canonical model, that is, without assuming exponential stratification. Inputs to the model are now examined.



Slide 7

A comparison of the canonical profiles of buoyancy frequency and potential sound speed gradient (thick solid lines) is made with averages obtained from STD casts. Significant differences occur, primarily in response to the Mediterranean Water intrusion.

In addition, a 17-meter rms internal wave amplitude was estimated from two thermistor arrays moored at 375 meters in the Azores Range. This compares with a 9-meter canonical amplitude for the same depth. Results of the non-canonical model are now shown.




Slide 8

Here, the non-canonical model is the internal wave μ^2 calculated using the actual ocean conditions. The canonical and the experimental curves are the canonical and directly-measured μ^2 profiles shown previously. Agreement between the non-canonical and experimental curves, particularly at the sound channel axis, suggests that the experimental μ^2 values result primarily from internal waves.

Furthermore, concurrent measurements in the Azores Range using towed, moored, and dropped sensors, plus intensive sound propagation experiments were also consistent with internal wave fluctuation statistics and with the high observed μ^2 levels.

MODEL VS. MEASUREMENT (MID-ATLANTIC RIDGE)



Slide 9

A similar set of sound speed profiles was collected at a second location, near the mid-Atlantic Ridge (~ 540 nautical miles SW of AFAR). The resulting direct, canonical, and non-canonical versions of μ^2 are shown. Again, use of the actual, rather than the idealized sound speed profile, can account for the large observed μ^2 values. The net result of these calculations is now shown.

COMPARISON OF VERY LOW FREQUENCY ATTENUATION
COEFFICIENTS FOR
DEEP SOUND CHANNEL, MID-LATITUDES, N. ATLANTIC

METHOD	ATTENUATION COEFFICIENT, α_1
CANONICAL MODEL	$0.5 \times 10^{-3} \text{dB/km}$
NON-CANONICAL MODEL	
AZORES RANGE	1.4
MID-ATLANTIC RIDGE	1.8
ATTENUATION MEASUREMENTS	1.3

Slide 10

Here the theoretical attenuation coefficient resulting from internal waves has been evaluated at the channel axis for both the Azores and the mid-Atlantic Ridge *experiment*. Also shown are attenuation coefficients based on the canonical model and on historical attenuation experiments in the North Atlantic. Slide 11 (not shown during the presentation) summarizes various parameter values used in the analysis. The predicted and measured coefficients agree closely when regional ocean conditions are used.

	GM 75	AZORES	M-AR
Z_1 (m)	1	1.5	1.3
N_1 (cph)	1.1	0.4	0.8
latitude	33°	$36^\circ 50'$	$28^\circ 18'$
n_1	0.37	0.13	0.27
n_1/w_i	24	8	20
μ_1^2	1.2×10^{-8}	1.5×10^{-7}	6.0×10^{-8}
j^*	3	3	3
B (km)	1	0.8	1.2
$\theta_0^2 \neq$.033	.022	.023
D_1	1.28×10^{-6}	1.87×10^{-6}	3.93×10^{-6}
α_1 (dB/km)	5.0×10^{-4}	1.4×10^{-3}	1.8×10^{-3}

\neq all θ_0^2 values standardized to a mean bottom depth of 3 km for N. Atlantic

EXPERIMENTAL α_1 FOR MID-LATITUDES IN N. ATLANTIC: 1.3×10^{-3} dB/km

Slide 11

In conclusion, we have presented evidence that appears to link the observed regional dependency of very-low frequency loss in the deep sound channel with an internal wave scattering mechanism. We are currently applying this methodology to data obtained in other areas.

Initial Distribution List

ADDRESSEE	NO. OF COPIES
NAVMAT (CAPT D. F. Parrish (MAT 08L), CAPT E. Young (MAT 0724))	2
NRL (B. Adams, R. Heitmeyer, A. Berman, Dr. K. M. Guthrie (Code 5160))	4
NOSC, San Diego (J. R. Lovett, Library)	2
NADC, Warminster	1
NCSL, Panama City	1
NPS, Monterey (Prof. C. N. Mooers, Prof. H. Medwin, Library)	3
NORDA, NSTL Station (R. Martin, R. Lauer, R. R. Goodman)	3
ONR (G. R. Hamilton, L. E. Hargrove (Code 421))	2
DARPA (CAPT V. P. Simmons)	1
NAVSEASYSKOM (C. D. Smith (Code 63R), R. W. Farwell (Code 63RA))	2
NUSC, Tudor Hill	1
COMNAVMARIANAS (CDR R. R. Miller)	1
University of Auckland (Prof. A. C. Kibblewhite)	1
Materials Research Laboratories (Dr. Daniel J. Whelan)	1
Deputy Adv. for Defence Science, Embassy of Australia (Dr. D. Wyllie)	1
Defence Scientific Establishment (Dr. Bert Olsson)	1
FWG (Dr. H.-G. Schneider)	1
Brown University, Dept. of Physics (Dr. Robert Beyer)	1
Catholic University of America, Dept. of Mechanical Engineering (Dr. Robert J. Urlick)	1
University of Houston, Cullen College of Engineering (Dr. Bill D. Cook)	1
Royal Australian Navy Research Laboratory (Dr. Martin Lawrence)	1
Admiralty of Underwater Weapons Establishment (Dr. David E. Weston)	1
Instituto de Pesquisas da Marinha (CDR Carlos Parente)	1
Yale University, Dept. of Engineering and Applied Science (Dr. John G. Zornig)	1
University of Tennessee, Dept. of Physics (Dr. Mack A. Breazeale)	1
Georgetown University, Physics Department (Dr. Walter G. Mayer)	1
Case Western Reserve University (Dr. Ernest B. Yeager)	1
Scripps Institute (Dr. F. H. Fisher, Dr. W. H. Munk)	2
University of Rhode Island, Dept. of Physics (Dr. Stephan Letcher)	1
APL/University of Texas (Dr. L. Hampton)	1
University of Bath, School of Physics (Dr. H. O. Berkday)	1
Defence Research Establishment, Atlantic (Dr. H. M. Merklinger)	1
Defence Research Establishment, Pacific (Dr. Robert Chapman)	1
University Louis Pasteur (Prof. S. Candau)	1
Technical university of Denmark (Dr. Leif Bjorno)	1
Defence Research Centre, Salisbury (Dr. Dennis H. Brown)	1
Laboratoire De Acoustica Submarina (Dr. Jorge Novarini)	1
DTIC	12

**DATE
FILMED
0-8**